
**U.S. Department
of Commerce**
Seattle, Washington

**Volume 101
Number 1
January 2003**

Fishery Bulletin

Contents

Articles

- 1–9** **Blaylock, Reginald B., Leo Margolis, and John C. Holmes**
The use of parasites in discriminating stocks of Pacific halibut
(*Hippoglossus stenolepis*) in the northeast Pacific
- 10–21** **Comyns, Bruce H., Richard F. Shaw, and
Joanne Lyczkowski-Shultz**
Small-scale spatial and temporal variability in growth and mortality of
fish larvae in the subtropical northcentral Gulf of Mexico: implications
for assessing recruitment success
- 22–31** **DeMartini, Edward E., Gerard T. DiNardo, and
Happy A. Williams**
Temporal changes in population density, fecundity, and egg size of
the Hawaiian spiny lobster (*Panulirus marginatus*) at Necker Bank,
Northwestern Hawaiian Islands
- 32–43** **Friedlander, Alan M., and David A. Ziemann**
Impact of hatchery releases on the recreational fishery for Pacific
threadfin (*Polydactylus sexfilis*) in Hawaii
- 44–57** **Hart, Deborah R.**
Yield- and biomass-per-recruit analysis for rotational fisheries, with an
application to the Atlantic sea scallop (*Placopecten magellanicus*)
- 58–74** **Hearn, William S., and Thomas Polacheck**
Estimating long-term growth-rate changes of southern bluefin tuna
(*Thunnus maccoyii*) from two periods of tag-return data
- 75–88** **Loefer, Joshua K., and George R. Sedberry**
Life history of the Atlantic sharpnose shark (*Rhizoprionodon*
terraenovae) (Richardson, 1836) off the southeastern United States

The conclusions and opinions expressed in *Fishery Bulletin* are solely those of the authors and do not represent the official position of the National Marine Fisheries Service (NOAA) or any other agency or institution.

The National Marine Fisheries Service (NMFS) does not approve, recommend, or endorse any proprietary product or proprietary material mentioned in this publication. No reference shall be made to NMFS, or to this publication furnished by NMFS, in any advertising or sales promotion which would indicate or imply that NMFS approves, recommends, or endorses any proprietary product or proprietary material mentioned herein, or which has as its purpose an intent to cause directly or indirectly the advertised product to be used or purchased because of this NMFS publication.

- 89–99** **Maunder, Mark N., and George M. Watters**
A general framework for integrating environmental time series into stock assessment models: model description, simulation testing, and example
- 100–115** **Miller, Michael J., David M. Nemerson, and Kenneth W. Able**
Seasonal distribution, abundance, and growth of young-of-the-year Atlantic croaker (*Micropogonias undulatus*) in Delaware Bay and adjacent marshes
- 116–128** **Newman, Stephen J., and Iain J. Dunk**
Age validation, growth, mortality, and additional population parameters of the goldband snapper (*Pristipomoides multidens*) off the Kimberley coast of northwestern Australia
- 129–146** **Ralston, Stephen, James R. Bence, Maxwell B. Eldridge, and William H. Lenarz**
An approach to estimating rockfish biomass based on larval production, with application to *Sebastes jordani*
- 147–167** **Winship, Arliss J., and Andrew W. Trites**
Prey consumption of Steller sea lions (*Eumetopias jubatus*) off Alaska: How much prey do they require?
- Notes*
- 168–174** **Beerkircher, Lawrence, Mahmood Shivji, and Enric Cortés**
A Monte Carlo demographic analysis of the silky shark (*Carcharhinus falciformis*): implications of gear selectivity
- 175–182** **Gunderson, Donald R., Mark Zimmermann, Daniel G. Nichol, and Katherine Pearson**
Indirect estimates of natural mortality rate for arrowtooth flounder (*Atheresthes stomias*) and darkblotched rockfish (*Sebastes crameri*)
- 183–188** **Marcogliese, David J., Elaine Albert, Pierre Gagnon, and Jean-Marie Sévigny**
Use of parasites in stock identification of the deepwater redfish (*Sebastes mentella*) in the Northwest Atlantic
- 189–193** **Polovina, Jeffrey J., Evan Howell, Denise M. Parker, and George H. Balazs**
Dive-depth distribution of loggerhead (*Caretta caretta*) and olive ridley (*Lepidochelys olivacea*) sea turtles in the central North Pacific: Might deep longline sets catch fewer turtles?
- 194–198** **Smith, Susan E., Robert A. Mitchell, and Dan Fuller**
Age-validation of a leopard shark (*Triakis semifasciata*) recaptured after 20 years
- 199** *Subscription form*

NOAA West Network

The file you requested could not be found.

You can either go to the [NOAA West Network](#) home page, or try our new [search engine](#).